Deflection, Spraying and Induced Scattering of Intense Laser Beams in Plasmas*

William L. Kruer

Lawrence Livermore National Laboratory University of California Livermore, CA 94550

Investigations into laser beam spraying, deflection, and induced scattering in plasmas are presented. Recent calculations and experiments on beam spraying¹ due to filamentation are discussed. A simple model is presented for the enhanced beam deflection ^{2,3} which occurs when the plasma flow transverse to the beam is nearly sonic. This model provides useful insights on the laser beam deflection, its scaling and the importance of self-consistent profile modifications. Finally, some initial discussion is given of recent experiments⁴ demonstrating the nonlinear behavior and competition of stimulated Raman and Brillouin scattering.

- 1. P. Young, J. Hammer, S. Wilks, and W. Kruer, Phys. Plasma 2, 2825 (1995).
- 2. Harvey A. Rose, Phys. Plasmas <u>3</u>, 1709 (1996); D.E. Hinkel et al., Phys. Rev. Letters (in press).
- 3. J. Moody, et al., submitted to Phys. Rev. Letters (1995); B. Bauer et al., to be published.
- 4. R. Kirkwood et al., J. Fernandez et al., and D. Montgomery et al., three papers submitted to Phys. Rev. Letters.

^{*}Work performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under Contract W-7405-ENG-48.